



Maxforce
PLATIN

BAYER **Maxforce**
PLATIN

REG. NO. L9986 Act No. 36 of 1947 **30 g**

INSECTICIDE

Cockroach

FOR FULL PARTICULARS SEE ENCLOSED PAMPHLET
Before using this product read the label carefully

An insecticidal gel bait for the control of American and German cockroaches in domestic premises, public hygiene and food storage premises.

ACTIVE INGREDIENT: Clothianidin (neonicotinoid).....10 g/kg

REGISTRATION HOLDER:
Bayer (Pty.) Ltd. Reg. No. 1968/011192/07
P. O. Box 143, ISANDO 1600,
SOUTH AFRICA, TEL: (011) 921 5911

MAXFORCE® is a registered trademark of the Bayer Group.

UN No. 3077 84907448A / 0219

CAUTION



Maxforce
PLATIN

INSECTICIDE

Reg. No. L9986 Act No. 36 of 1947

Before using this product read the label carefully.

An insecticidal gel bait for the control of American and German cockroaches in domestic premises, public hygiene and food storage premises.

ACTIVE INGREDIENT: Clothianidin (neonicotinoid).....10 g/kg

REGISTRATION HOLDER: Bayer (Pty.) Ltd.

Reg. No. 1968/011192/07
P.O. Box 143, ISANDO 1600, SOUTH AFRICA
TEL: (011) 921 5911

MAXFORCE® is a registered trademark of the Bayer Group.

4 x 30 g

84923567A / 0219



Cockroach



CAUTION
VERSIGTIG



WARNINGS:

Handle with care.
Harmful when swallowed.
Harmful to fish and other animals.
Store in a cool place and away from food and feedstuffs.
Keep out of the reach of children, uninformed persons and animals.

PRECAUTIONS:

Avoid contact by mouth and skin.
Wear rubber gloves during application.
Wash with soap and water immediately after use or accidental skin contact.
Do not eat, drink or smoke whilst applying and before washing hands and face.
Apply only in positions where it will be completely inaccessible to children and pets.
Do not contaminate dams, drains, ponds, streams and other waterways with the product or used containers.
Do not apply gel to surfaces on which food or feed is stored, prepared or eaten.
Protect food preparing equipment and eating utensils from contamination during application.
Destroy used container and do not re-use for any other purpose.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

For maximum effectiveness **Maxforce® Platin** must be placed in and around cockroach harbourages and, if possible between these areas and food sources. Survey the area to be treated to identify sites of cockroach infestation e.g. by using traps and/or flush-out aerosols, searching for droppings, cast skins and egg cases etc.

Maxforce® Platin can be placed in the vicinity of: corners, cracks and crevices, voids, service-ducts, lift shafts, equipment and furniture e.g. counters, refrigerators, cookers, sinks, baths etc. **Maxforce® Platin** is particularly valuable in sensitive areas where the use of conventional insecticide formulations is restricted e.g. in the vicinity of electric and electronic equipment etc. Where immigration of cockroaches has been identified placement of additional bait in peripheral areas of the site will aid control.

Apply **Maxforce® Platin** directly as spots or thin ribbons to surfaces in identified target areas. It should be placed out of sight and where light intensity is low. Place the bait spots in places inaccessible for children or pets.

Avoid application to excessively dusty, damp or greasy locations, and areas which are subject to washing.
Do not apply to areas that have recently been treated with other insecticides or contaminate the bait with other insecticides.

Maximum levels of control are achieved between six days and several months after treatment depending upon the level of infestation and as long as the gel is present. Where infestations are high inspect the applied spots of **Maxforce® Platin** regularly and make further applications as required.

Maxforce® Platin 30 g is dispensed using a purpose-built caulking gun, which is used as follows: Insert the cartridge into the gun frame. Remove the cap of the cartridge tip. Insert a dispensing tip as supplied with the caulking gun. A plastic dispensing tip should be used in and around electrical installations and equipment. Press the trigger of the caulking gun until the plunger contacts the piston of the cartridge. The next application of pressure to the trigger should release a droplet of gel from the dispensing tip.

Application rates:

Bait application should equal approximately 0,25-0,5 g / spot and 10 to 20 spots / 10 m².

- A spot of approximately 3 mm diameter equates to 0,25 g of bait.
- A spot of approximately 6 mm diameter equates to 0,5 g of bait.
- Low infestation: cockroaches rarely visible during the day.
- High infestation: cockroaches commonly visible during the day.

The lower application rate should be used for low and moderate German cockroach infestations or re-treatments. The higher rate should be used for more severe infestations of American and German cockroaches or first treatment.

Numerous smaller placements will provide faster control than a few larger spots. The gel will remain pliable and palatable to cockroaches as long as it is visible; usually up to 12 weeks. Where infestations are high inspect gel four weeks after treatment and replace if necessary. The life of the bait will reduce under daylight conditions.

Maxforce® Platin can be integrated with space or surface spray treatments but must not be contaminated with other insecticides or placed on freshly sprayed surfaces.

For any product enquiry, contact Bayer Environmental Science - 011 921 5911. For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

© Bayer (Pty.) Ltd. All unauthorized reproduction of this label is forbidden.
www.environmentalscience.bayer.co.za

WAARSKUWINGS:

Hanteer versigtig.
Skadelik indien ingesluk.
Skadelik vir vis en ander diere .
Bewaar in 'n koel, droë plek weg van voedsel en voer.
Hou buite die bereik van kinders, diere en oningeligte persone.

VOORSORGMATREËLS:

Vermý alle kontak deur mond of met vel.
Dra rubber handskoende tydens aanwending.
Was met seep en water onmiddelik na gebruik of toevallige velkontak.
Moet nie eet, drink of rook tydens aanwending nie of voordat gesig en hande gewas word nie
Wend slegs aan in posisies wat ontoeganklik is vir kinders en troeteldiere.
Moet nie damme, dreine, poele, strome of enige ander waterweë kontamineer met die produk of leë produk houers nie.
Moet nie die jel toedien op oppervlaktes waarop kos of voer geberg, voorberei of geëet word nie.
Voorkom besoedeling van voedselvoorbereiding toerusting en eetgerei tydens aanwending Vernietig leë houers en moet dit nie vir enige ander doeleindes gebruik nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI.

Vir optimale resultate moet **Maxforce® Platin** binne-in en rondom kakkerlak skuilplekke aangewend word, asook tussen hierdie areas en voedselbronne. Ondersoek die area wat behandel moet word om enige areas van kakkerlak infestaties te identifiseer bv. gebruik lokvalle en / of uit-spoel aërosols, soek vir mis, doppe en eier pakkies ens.

Maxforce® Platin kan geplaas word in / naby: hoeke, krake, skeure, gapings, diens kanale, huisbak skagte, toestelle en meubels bv. toonbanke, yskaste, oonde, wasbakke, baddens ens. **Maxforce® Platin** is veral waardevol in sensitiewe areas waar die gebruik van konvensionele insekdoder formulasies beperk is bv. in die omgewing van elektriese en elektroniese toestelle ens. Waar immigrasie van kakkerlakke waargeneem is, sal die addisionele plasing van lokaas in die omliggende areas help met die beheer van die kakkerlakke.

Maxforce® Platin kan as 'n kolbehandeling of dun stroke aangewend word op geïdentifiseerde teiken areas, verkieslik buite sig en waar die ligintensiteit laag is. Plaas die kolle in areas wat ontoeganklik is vir kinders of troeteldiere.

Vermý oormatige stowwerige, klam of vetterige plekke.
Vermý applikasie op oppervlaktes wat onlangs behandel is met ander insekdoders of kontaminasie van die lokaas met ander insekdoders.

Maximale vlakke van beheer word bereik tussen 6 dae en 'n paar maande na behandeling, afhange van die vlakke van infestatie en hoe lank die jel teenwoordig is. Waar infestaties hoog is, inspekteer die toegediende kolle van **Maxforce® Platin** gereeld en maak addisionele applikasies indien nodig.

Maxforce® Platin 30 g is toegedien deur gebruik te maak van 'n doel-geboude “caulking” geweer, wat as volg gebruik word: Plaas die “patroon” binne-in die geweer raam. Verwyder die doppe vanaf die “patroon” tip. Voeg die resepterende tip by, wat verskaf word saam met die “caulking” geweer. 'n Plastiese resepterende tip moet gebruik word binne-in en rondom elektriese installasies en – toestelle. Druk die sneller van die “caulking” geweer totdat die druppunt in kontak is met die suier van die patroon. Die volgende toepassing van druk op die sneller moet 'n druppel jel vrylaat vanuit die resepterende punt.

Dosis vir toediening:

Lokaas toediening moet ongeveer gelyk wees aan 0.25 – 0.5g / kol en 10 – 20 s / 10 m²

- 'n Kol van ongeveer 3 mm in deursnee is gelyk aan 0.25g lokaas.
- 'n Kol van ongeveer 6 mm in deursnee is gelyk aan 0.5g lokaas.

- Lae infestatie: Kakkerlakke skaars sigbaar gedurende die dag.
- Hoë infestatie: Kakkerlakke algemeen sigbaar gedurende die dag.

Die laer dosis van toediening moet gebruik word vir lae en matige Duitse kakkerlak infestaties of vir herbehandelinge. Die hoër dosis moet gebruik word vir meer ernstige infestaties van Amerikaanse en Duitse kakkerlakke of eerste behandelinge.

Veelvoudige kleiner plasing sal vinniger beheer verskaf as 'n paar groter kolle. Die jel sal buigsaam en aanvaarbaar wees vir kakkerlakke so lank as wat dit sigbaar is; gewoonlik tot in met 12 weke. Waar infestaties hoog is, ondersoek die jel vier weke na behandeling en vervang indien nodig. Die lewe van die lokaas sal verminder met daglig omstandighede.

Maxforce® Platin kan geïntegreer word met area of oppervlak sproei behandelinge, maar moet nie gekontamineer word met ander insekdoders of, geplaas word op vars bespuite oppervlaktes nie.

Vir enige produk navrae, skakel Bayer Environmental Science - 011 921 5911. Slegs vir vergiftigingsnoodgevallen kan Griffon Gif Inligtingentrum ook gekontak word - 082 446 8946.

© Bayer (Edms.) Bpk. Alle onregmatige kopiëring van die etiket is verbode.
www.environmentalscience.bayer.co.za