



**Racumin<sup>®</sup> 3D**  
PASTE



# Racumin<sup>®</sup> 3D

## PASTE

### RODENTICIDE



REG. NO. L10218 Act No. 36 of 1947

Before using this product read the label carefully.

**A universal anticoagulant bait for the control of rats and mice indoors and outdoors.**

**ACTIVE INGREDIENT:**

Coumatetralyl (anti-coagulant).....0,375 g/kg

**CO-FORMULANT:**

Cholecalciferol.....0,10 g/kg

**REGISTRATION HOLDER:**


Bayer (Pty.) Ltd.  
Reg. No. 1968/011192/07  
P.O. Box 143, ISANDO 1600, SOUTH AFRICA  
TEL: (011) 921 5911

RACUMIN<sup>®</sup> is a registered trademark of the Bayer Group.

UN No. 3027



Rats • Mice



**5 kg e**

ZA85829181D - ARTICLE 85840916



CAUTION  
VERSIGTIG





**Racumin<sup>®</sup> 3D**  
**PASTE**

ZA85837095C - ARTICLE 85840916



**Racumin<sup>®</sup> 3D**  
**PASTE**

**WARNINGS:** Avoid contact with skin, eyes and mouth. May cause sensitization by skin contact. Handle with care. Harmful to fish and wildlife.

**PRECAUTIONS:** Keep containers tightly closed in a dry, cool and well-ventilated place. Keep away from food, drink and animal feedingstuffs. Store in original container. Store in a place accessible by authorized persons only. Store unused sachets in a safe place. DO NOT STORE parts of used sachets. Wear rubber gloves during application.

**DIRECTIONS FOR USE:** USE ONLY AS DIRECTED.

• **For outdoor and indoor uses where children or non-target organisms could be present:**

The bait sachets must be placed in a tamper-proof or lockable, secured bait stations, or placed, secured with wire or nail and covered, in a manner to be inaccessible to children and non-target organisms.

• **For indoor uses where no children or non-target organisms could be present:**

The baits can be placed and fixed in a bait station or applied loose in hidden places. Inspect baiting at least in weekly intervals, and refill according to consumption. It is important to provide surplus of bait during the first days of the treatment. Rat and mouse control measures should continue until no more bait consumption is apparent, but not exceeding five weeks. A normal bait campaign lasts 2 to 3 weeks. Spoiled bait and carcasses have to be collected during every inspection of the site. When rodent control is achieved, all remaining baits must be collected and disposed of as pesticidal waste.

**APPLICATION RATES:**

PESTS	DOSAGE RATE	APPLICATION
General Public Health		
<b>RATS</b>	100g to 200g	Place <b>RACUMIN 3D Paste</b> per baiting point depending upon the infestation level (from low to high infestation, respectively). An amount of 200g of product must be added when bait has been totally consumed within one control interval. Subsequent follow up visits must be no more than 14 days apart. <b>Distance between baiting points:</b> 5-20 linear metres for a low infestation. 3-10 linear metres for a heavy infestation.
<b>MICE</b>	10g to 20g	Place <b>RACUMIN 3D Paste</b> per baiting point depending upon the infestation level (from low to high infestation, respectively). An amount of 20g of product must be added when bait has been totally consumed within one control interval. Subsequent follow up visits must be no more than 14 days apart. <b>Distance between baiting points:</b> 5-20 linear metres for a low infestation. 2-10 linear metres for a heavy infestation.

Application is recommended on sites either where mice or rats feed or on the rat runways. Baiting points are to be inspected weekly, and bait to be replenished according to consumption. Due to its high fat-content, no application is recommended in places, which may be exposed to direct sunlight.

For any enquiry, contact Bayer - **0861 022 937 (08610BAYER)**. For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

© Bayer (Pty.) Ltd. All unauthorized reproduction of this label is forbidden.

Batch no:  
Date of manufacture:

- Coumatetralyl
- Cholecalciferol

Content: Coumatetralyl 0.0375 %; Cholecalciferol 0,01%

Formulation: Bait (ready for use) (RB)

**WARNING**



WHO-classification: III (Slightly hazardous)

**WARNING**

**SYMPTOMS OF HUMAN POISONING:**

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around the elbow and knee joints and blood in urine and faeces.

**FIRST AID TREATMENT:**

If accidentally swallowed contact a physician immediately.

**NOTE TO PHYSICIAN:**

Intravenous Vitamin K1, 50-150 mg on first day depending on prothrombin time. In severe cases small transfusions of fresh blood.

**PRECAUTIONARY STATEMENTS**

P280 Wear protective gloves/protective clothing/eye protection/face protection.  
P333 + P313 If skin irritation or rash occurs: Get medical advice/attention.  
P501 Dispose of contents/container in accordance with local regulation.

**OTHER HAZARDS**  
No other hazards known.



6 005317 001201

**Bayer**